ENGLISH – PATIENT ADVICE SHEET If you forget to take your progestogen-only pill (mini-pill)

What is the time limit for a missed pill?

It depends on the type of progestogen-only pill you are using. Here are the main situations:

- Pill containing **desogestrel 75μg** (Cerazette®, Optimizette®, Antigone®, or generics) → tolerated delay of 12 hours.
- Pill containing **levonorgestrel 30μg** (Microval®) → tolerated delay of **3 hours**.
- Pill containing drospirenone 4 mg (Slinda®) → tolerated delay of 24 hours, but the instructions differ. → Check the leaflet.
- [2] If in doubt, check the name on the box or prescription, or ask a healthcare professional.

What to do

✓ If the delay is equal to or less than the tolerated time:

- Take the missed pill as soon as possible.
- Take the next pill at the usual time.
- → Contraceptive effectiveness is maintained.
- → No additional precautions are needed.

X If the delay exceeds the tolerated time:

• Take the missed pill immediately, even if two pills are taken on the same day or at the same

If more than one pill was missed, discard the others to avoid mistakes.

- ▲ If unprotected sex occurred in the past 5 days → emergency contraception is recommended.
- Take a urine pregnancy test 3 weeks after the at-risk intercourse, even if emergency contraception was used.
- Use a condom for the following 7 days.
- Continue the pack as normal.

Emergency contraception

- Morning-after pill:
 - Norlevo®: up to 72h
 - EllaOne®: up to 120h
 - Copper IUD: up to 5 days after unprotected sex

Helpful tips

- There is never a break between packs.
- Take the pill at the same time every day.
- Vomiting or severe diarrhea < 4h after taking = equivalent to a missed pill.
- Use an alarm or reminder app.
- If in doubt, ask your doctor, midwife, or pharmacist.